

Manufacturer's Information Test No. 20-5002: Drug Analysis

Each sample pack consisted of two items. Item 1 contained 300 mg of powder consisting of corn starch. Item 2 contained 300 mg of powder consisting of approximately 3.3% oxycodone hydrochloride and 96.7% lactose monohydrate. Oxycodone hydrochloride is a Schedule II controlled substance in the United States.

SAMPLE PREPARATION-

The corn starch in Item 1 was a pure powder. The oxycodone hydrochloride in Item 2 was a pure powder that was combined with lactose monohydrate. These two powders were mixed thoroughly to ensure homogeneity.

ITEM 1 (PREPARATION): Approximately 300 mg of the corn starch powder was weighed out and deposited into a glassine bag which was folded and secured with a label. The folded glassine bag was placed into a small zip top bag, which was heat sealed and then placed into a pre-labeled envelope.

ITEM 2 (PREPARATION): Approximately 300 mg of the powder containing oxycodone hydrochloride and lactose monohydrate was weighed out and deposited into a glassine bag which was folded and secured with a label. The folded glassine bag was placed into a small zip top bag, which was heat sealed and then placed into a pre-labeled envelope.

SAMPLE PACK ASSEMBLY: One of each of the Item 1 and Item 2 envelopes was placed into a larger pre-labeled sample pack envelope.

VERIFICATION: The above substances were identified by predistribution laboratories, who used the following combined list of techniques: color tests, UV, and GC/MS.

The information presented here details how test samples were prepared as well as any design specifications. This information does not necessarily represent the answers that should or could be obtained from an examination of the sample(s). Final interpretation of the results should be deferred until the summary report is available.